

Seasonality in Extreme Precipitation as Depicted in the TMPA and GPCP 1DD

George J. Huffman^{1,2,4}, Robert F. Adler^{1,3}, David T. Bolvin^{1,2}, Eric J. Nelkin^{1,2}
 1: NASA/GSFC Mesoscale Atmospheric Processes Laboratory
 2: Science Systems and Applications, Inc.
 3: Univ. of Maryland College Park/ESSIC
 4: george.j.huffman@nasa.gov
<http://precip.gsfc.nasa.gov>

1. Introduction

It is interesting to compute extremes in satellite data sets in order to
 - gain an understanding of the climatology of the metrics
 - gain an understanding of the satellite data set limitations

The precipitation group in the NASA/GSFC MAPL is responsible for two high-resolution precipitation products
 - TRMM Multi-satellite Precipitation Analysis (TMPA)
 - GPCP One-Degree Daily (1DD)

Neither data set is specifically designed for extremes

There is also a scale mismatch
 - TMPA is averaged to UTC days, since that is the conventional minimum period for extremes, and matches the 1DD time
 - TMPA is computed at both 0.25° and 1° resolution for information about scale dependence

As well, 1DD provides information at high latitudes

Seasonal behavior is a key part of the climate signal

The globe is broken into regions for easier viewing

2. TRMM Multi-satellite Precipitation Analysis (TMPA)

3-hr 0.25° grid, 50°N-S

Microwave precip:

- intercalibrate, combine

IR precip:

- calibrate with microwave


Combined microwave/IR:

- IR fills gaps in microwave

Final:

- accumulate combined 3-hr precip for the month
 - weighted combo. with gauge analysis
 - rescale 3-hr precip to sum to the monthly sat-gauge combination

Real-time version faded out



3. One-Degree Daily (1DD)

1-day, 1° grid, 90°N-S

40°N-S:


- Threshold-Matched Precip Index (TMPI)
 - month of coincident GPROF-SSM/I precip, geo-IR Tb data to set precip fraction (at 1° resolution) --> local IR Tb threshold
 - local conditional rate from GPCP monthly
 - Like GPI with IR threshold, single conditional rate varying by month, location

Outside 40°N-S:

- daily TOVS/AIRS precip occurrence scaled to TMPI occurrence at 40°N-S
 - daily TOVS/AIRS rates rescaled to sum to GPCP monthly locally

Final:

- local linear fade from TMPI to rescaled TOVS/AIRS precip by day, 40-50°



4. Data Sources

A diverse, changing set of input precip estimates – various

- periods of record
 - regions of coverage
 - sensor-specific strengths and limitations


TMPA designed to give the “best” instantaneous estimate

- input data sources vary
 - short dashes indicate data being used in the new Version 7

- monthly TMPA SG
 - 0.25°, 50°N-S, 3-hr→daily

1DD more approximate, relatively homogeneous

- match-up of single SSM/I (soon SSMIS), geo-IR
 - monthly GPCP SG
 - 1°, 90°N-S, daily



5. Climate-oriented Indices

Acknowledge CCI/CLIVAR/JCOMM Expert Team (ET) on Climate Change Detection and Indices (ETCCDI) concept of “core indices”

Chose to compute

- Ravg Avg. daily precip (=PTOT/365)
- Rfrac Avg. fraction of days with precip (> 0.5 mm/d)
- R95p 95th-percentile precip rate
- CDD Avg. annual maximum length of dry spell (<1 mm/d)
- CWD Avg. annual maximum length of wet spell (≥1 mm/d)

Record is too short to compute sophisticated metrics!

Note the paradox of “climate” variables depending on fine-scale estimates

- “extremes” easily contaminated by analysis artifacts
- R95p is computed because it is well-correlated to 99th percentile and maximum values, and is more stable

Introduce a dryness index:

- f2mm Avg. fraction of days with precip ≤ 2 mm/d
- rough lower limit of agriculturally relevant event
- less sensitive to analysis artifacts than CDD

6. Results – Seasonal Ravg


Minimal scale dependence

Algorithm similarities

- Southern Ocean minimum DJF, maximum MAM-JJA
- Tasman Sea, New Zealand minimum DJF, maximum JJA
- expected summer monsoon cycle in S. Asia
- break in ITCZ off C. America JJA-SON
- double Pac. ITCZ MAM

Algorithm differences

- Tasmania, New Zealand much wetter in 1DD
- N. Pac. storm track wetter in 1DD JJA-SON
- Patagonia and offshore wetter in 1DD
- low-precip artifact off Newfoundland in TMPA



7. Results – Seasonal f2mm


Moderate-strong scale dependence (note Borneo)

Algorithm similarities

- Southern Ocean minimum MAM, maximum SON-DJF, leads Ravg
- eastern Indian Ocean has a decent seasonal cycle, unlike Ravg
- strong seasonal cycle in central Africa, Asian monsoon regions
- minimum offshore of N.E. Brazil JJA when onshore region has a maximum

Algorithm differences

- 3B42 higher in Southern Ocean
- Tasmania and New Zealand stick out in 3B42, don't in 1DD



8. Results – Seasonal R95

Strong scale dependence

Algorithm similarities

- peaks tend to be in phase with Ravg
- peaks around Darwin, offshore of S.E. Asia and the Philippines, southwest of C. America are unlike Ravg
- strong maxima in la Plata basin, S. America; note seasonal shift in location

Algorithm differences

- Tasmania, New Zealand much wetter in 1DD, as with Ravg
- 1DD tends to higher minima, particularly in subtropical highs
- bigger break across S. Andes in 3B42, except DJF
- 1DD has a break at 40°S, less so at 40°N

