

Impact of the Initialization Method on the Skill of Decadal Climate Predictions in the Southern Ocean

UCL **Violette Zunz**¹ (violette.zunz@uclouvain.be), Hugues Goosse¹, Johann Jungclaus², Svetlana Dubinkina¹ and Yoann Sallaz-Damaz¹

¹Georges Lemaître Centre for Earth and Climate Research, Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium.

²Max Planck Institute for Meteorology, Hamburg, Germany.

1. Introduction

Performing efficient decadal climate predictions over the Southern Ocean relies partly on:

- a model that represents relatively well the physical process in this region;
- an optimal estimate of the initial state of the system.


Data assimilation (DA) methods can provide an optimal estimate of the initial state given imperfect model equations and inaccurate and incomplete observations.

Objective: testing a sophisticated DA method to check if it can improve decadal climate predictions skill in the Southern Ocean area.

2. Methodology

- ### 2.1. Model
- Coupled climate model **LOVECLIM** [1]:
- 3D Earth system model of intermediate complexity;
 - Lower level of complexity and coarser resolution than general circulation model (GCM);
 - Lower computational cost than GCM.
- ➔ Large number of tests can be performed within a reasonable computational time.

- ### 2.2. Data Assimilation: Combination of a Particle Filter and Nudging
- Used to reconstruct the evolution of the climate system between 1850 and 2010;
 - Variable assimilated: anomaly of the surface temperature w.r.t. 1961-1990 from HadCRUT3 dataset [2].




- **Nudging**
- $k(T_{mod} - T_{obs})$
is added to the heat flux between the atmosphere and the ocean (for each grid point of the ocean free of sea ice), in order to nudge the solution towards the observation.
 T_{mod} and T_{obs} = modeled and observed surface temperature anomaly;
 k = relaxation coefficient ($\sim$ relaxation time of 6 months).
Nudging limited to a maximum flux of 50 Wm^{-2} .


- ### 2.3. Hindcast Experiments
- 5-year ensemble simulations (96 members):
 - Start in January, every 2 years from 1980 to 2002;
 - 10-year ensemble simulations (96 members):
 - Start in January, every 5 years from 1975 to 1990;
 - 2 series of hindcasts for each time period: one series whose initial condition has been extracted after the resampling step of the particle filter (**hindcasts with DA**), the other one initialized without DA (**hindcasts without DA**).

The accuracy of these hindcasts is assessed by comparing their results with available observations. Hindcasts performed with LOVECLIM are compared to the ones from the ECHAM6/MPI-OM model in section 5.

3. Reconstruction of the Surface Temperature




4. Reconstruction of the Trends of the Sea Ice Concentration



5. 10-Year Hindcasts Performed with LOVECLIM and with ECHAM6/MPI-OM

Hindcasts performed with the **ECHAM6/MPI-OM** atmosphere-ocean-sea-ice coupled model are **10 members** ensemble. They are initialized through the assimilation (with only a nudging) of the 3D temperature and salinity anomalies (except in the area covered by sea ice) taken from an ensemble of MPI-OM ocean experiments forced with the NCEP-NCAR atmospheric forcing [6].



References

[1] Goosse H. et al., 2010. *Description of the Earth system model of intermediate complexity LOVECLIM version 1.2*, Geosci. Model Dev., 3, 603-633, 10.5194/gmd-3-603-2010.

[2] Brohan P. et al., 2005. *Uncertainty estimates in regional and global observed temperature changes: a new dataset from 1850*, J. Geophys. Res., 111, D12106, 10.1029/2005JD006548.

[3] Dubinkina S. et al., 2011. *Testing a particle filter to reconstruct climate changes over the past centuries*, Int. J. Bifur. Chaos (in press).

[4] van Leeuwen, P. J., 2009. *Particle Filtering in Geophysical Systems*. Monthly Weather Review, 137(12):4089-4114. 10.1175/2009MWR2835.1.

[5] Comiso, J., 1999, updated 2008. *Bootstrap Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I, 1979-2007*. Boulder, Colorado USA: National Snow and Ice Data Center. Digital media.

[6] Matei, D. et al., 2011. *Two tales of initializing decadal climate predictions experiments with the ECHAM5/MPI-OM model*, Journal of Climate (to be submitted).

[7] Wilks D. S., 2006. *Statistical Methods in the Atmospheric Sciences*, 2nd edition, Academic Press, 627 pp.

6. 5-Year Hindcasts Performed with LOVECLIM

6.1. Accuracy Measure of the Hindcasts [7]

Proportion Correct (PC):
 $PC = \frac{a+d}{n}$; $n = a+b+c+d$.
PC = 1 for perfectly accurate hindcasts.

Hindcast	Case 1	Observed	
		Case 1	Case 2
	Case 1	a	b
	Case 2	c	d

6.2. Proportion Correct of 5-Year Trends of Sea Ice Extent and Area

case 1 $\rightarrow$ trends > 0 ; case 2 $\rightarrow$ trends < 0 .

6.3. Proportion Correct of 5-Year Trends of Sea Ice Concentration


6.4. Proportion Correct of 5-Year Trends of Sea Ice Extent Dipole between Bellingshausen/Amundsen and Ross Sectors

For some time period, observed 5-year trends of sea ice extent presents a dipole between Bellingshausen/Amudsen Seas and Ross Sea, i.e. if the sea ice extent trends in one area is positive, it is negative in the other area and conversely.

case 1
 $\rightarrow$ Ross > 0 and Bellingshausen/Amundsen < 0 ;
case 2
 $\rightarrow$ Ross < 0 and Bellingshausen/Amundsen > 0 .

	Hindcasts with DA		Hindcasts without DA	
	Extent	Area	Extent	Area
Weddell Sea	0,38	0,25	0,50	0,50
Indian Ocean	0,33	0,22	0,11	0,22
Western Pacific Ocean	0,43	0,43	0,00	0,29
Ross Sea	0,57	0,43	0,14	0,14
Bellingshausen/Amundsen Seas	0,75	0,75	0,75	0,75
Southern Ocean	0,11	0,11	0,22	0,22

Hindcasts with DA	Hindcasts without DA
0,50	0,38



7. Discussion

- Data assimilation of surface temperature anomaly clearly improves the reconstruction of the surface temperature in the Southern Ocean area.
- Sea ice concentration trends between 1979 and 2007 computed by the model with DA present pattern closer to the observed trends than the model without DA.
- Caution must be taken while initializing hindcasts in the Southern Ocean: hindcasts initialized with the nudging can present non physical oscillations of the sea ice area in several sectors of the Southern Ocean, these oscillations appearing later in the simulation with LOVECLIM than with ECHAM6/MPI-OM.
- 5-year hindcasts performed with LOVECLIM shows that even if the proportion correct is not very high for both hindcasts initialized with and without DA, the score is usually better for hindcasts initialised with DA, especially in the Ross Sea sector.
- In conclusion, data assimilation of the surface temperature used to initialize hindcasts seems to improve slightly the forecast skill of the Southern Ocean sea ice at decadal time scale.

8. Next Steps

- Deeper investigation of the behaviour of Southern Ocean sea ice in hindcasts initialized with DA.
- Test of an initialization method wich doesn't trigger non physical oscillation of the Southern Ocean sea ice.
- Improvement of the combination of the particle filter with the nudging by taking into account the nudging in the computation of the weight of the particles.
- Tests with assimilation of 3D temperature and salinity in the ocean.
- Implementation of a method to assess the quality of hindcasts of Southern Ocean sea ice based on Empirical Orthogonal Function (EOF) to point out spatial mode of variability.