


# Using "Replay" to Data Assimilation for AGCM Model Development

Andrea Molod  
Lawrence Takacs  
Tommy Owens

NASA/GMAO



NASA Goddard Space Flight Center  
GLOBAL MODELING AND ASSIMILATION OFFICE



4<sup>th</sup> World Climate Research Programme  
International Conference on  
Reanalyses

7-11 May 2012  
Silver Spring, Maryland, USA






# Outline

- Example where data assimilation experiments at  $0.25^\circ$  exhibited undesired behavior in response to an AGCM change which improved simulated climate
- “Replay” experiments at  $1^\circ$  used as a proxy for running data assimilation experiments to explore parameter sensitivity with new model
- Confirmation of “replay” results with data assimilation experiments




# Issue: Impact of AGCM Change in Moist Processes

Data Assimilation at  $0.25^\circ$  -> Precipitation changes with model change



Analysis  
Moisture  
Increment  
(g/kg/day)



**GOAL: Use new model in Data Assimilation mode with better parameter choices**

# Issue: Impact of AGCM Change in Moist Processes

\* Issue traced to change in model parameters to describe the sub-grid scale probability distribution of total water. Width of PDF ~ AGCM "Critical Relative Humidity" (RHcrit).

\* Smaller RHcrit in new AGCM results in:

More precipitation and larger analysis increment (Humidity is constrained)


Strategy --> Explore parameter space for data assimilation mode. Seek out smaller analysis increment and correct total precipitation

Approach --> Run "Replay" experiments at 1°







NASA Goddard Space Flight Center  
GLOBAL MODELING AND ASSIMILATION OFFICE





# "Replay" at GMAO: Flow Diagram







# Comparison of 1° Replay to 0.25° Assimilation



Temperature



Specific Humidity




Precipitation

Replay reproduces assimilation analyzed fields, approximates assimilation derived fields




NASA Goddard Space Flight Center  
GLOBAL MODELING AND ASSIMILATION OFFICE



# Replay with different RHcrit - Total Precipitation




**RHcrit: 0.80**



**RHcrit from Original AGCM**



**RHcrit: 0.90**





**RHcrit: 0.95**

Replay experiments were performed with RHcrit ranging from 0.8 to 0.98. Precipitation similar to original AGCM result when RHcrit=0.95





NASA Goddard Space Flight Center  
GLOBAL MODELING AND ASSIMILATION OFFICE

# Replay with different RHcrit Values




Replay data increment of moisture in g/kg/day




Data increment similar to original AGCM results when RHcrit=0.95



# Confirmation of "Replay" result: Data Assimilation Experiments at 0.5°



**RHcrit: 0.95**



**"Original AGCM"**






Full data assimilation experiment with RHcrit=0.95 show increments and precipitation resembling the result using the original model formulation. Replay result is borne out.



NASA Goddard Space Flight Center  
GLOBAL MODELING AND ASSIMILATION OFFICE

# Lessons learned for AGCM Development: Dependance on Horizontal Resolution

Before: GEOS-5 AGCM Specific Humidity is sensitive to resolution






$2^\circ \times 2.5^\circ$

$1^\circ \times 1^\circ$

$0.5^\circ \times 0.67^\circ$

GEOS-5 AGCM Total Precipitation is insensitive to resolution



Initial parameter choice based on coarse resolution AGCM  $\rightarrow$  DAS precipitation sensitivity and resulting parameter choice revealed resolution dependance




NASA Goddard Space Flight Center  
GLOBAL MODELING AND ASSIMILATION OFFICE

# RHcrit Dependence on Horizontal Resolution

Total water field from a GEOS-5 simulation at ~10 km resolution on the cubed sphere was regrided to coarser resolution, and sub-grid variance was computed.

Sub-grid scale variance of total humidity is related to "Critical relative humidity" used for moist processes.




General vertical profile of the critical relative humidity is unchanged as the grid resolution changes, for all regions and underlying surfaces, and shifts to higher values with smaller grid size.



# Dependance on Horizontal Resolution

After: GEOS-5 AGCM with resolution dependant RHcrit



# Summary

- “Replay” experiments exhibit DAS behavior at the resolution of the constraining assimilation. That is, replay experiments at  $1^\circ$  reflect the behavior of the  $0.25^\circ$  analysis.
- Data assimilation experiments at  $0.5^\circ$  confirmed the results of replay experiments --> Replay can be used to assess behavior in DAS mode
- Results from examining DAS sensitivity used to improve AGCM at high resolution

