

## Table of Contents

### 1. Atmospheric data assimilation schemes, analysis and initialization, data impact studies, observing system experiments

| Author  | Paper Title  | Country | Pages |
|---|--|---------|-------|
| K. Araki, M. Murakami, T. Kato, T. Tajiri   | Diurnal variation of thermodynamic environments for convective cloud development around the central mountains in Japan during warm seasons | Japan   | 1-03  |
| K. Araki, M. Murakami, H. Ishimoto, T. Tajiri   | The 1-dimensional variational approach to improve thermodynamic profiles in low-level troposphere during rain conditions                   | Japan   | 1-05  |
| K. Araki, M. Murakami   | Validation of vertical thermodynamic profiles by cloud base temperature obtained from a ground-based infrared radiometer                   | Japan   | 1-07  |
| J. Fukuda, T. Tsukamoto, T. Fujita  | Improvement of JMA's meso-scale analysis background errors   | Japan   | 1-09  |
| Y. Ikuta  | Data assimilation using GPM/DPR at JMA   | Japan   | 1-11  |
| M. Kazumori   | Assimilation of GPM microwave imager data in JMA's NWP systems   | Japan   | 1-13  |
| M. Kazumori   | Assimilation of Himawari-8 clear sky radiance data in JMA's NWP systems  | Japan   | 1-15  |
| M. Kazumori   | Assimilation of Megha-Tropiques/SAPHIR radiance data in JMA's global NWP system  | Japan   | 1-17  |
| E. Klimova  | Stochastic ensemble Kalman filter with ensemble generation using a square root form: ensemble $\pi$ -algorithm                             | Russia  | 1-19  |
| Y. Kosaka   | Assimilation of surface pressure in METAR  | Japan   | 1-21  |
| M. Moriya   | Operational use of ASCAT ocean vector wind data in JMA's mesoscale NWP system  | Japan   | 1-23  |
| M. Pondaca, S. Levine, R. Yang, A. Gibbs, J. Carley, G. Manikin, C. Gaustini, Y. Lin, J. Purser, X. Su, J. Whiting, D. Parrish, G. DiMego | Development progress on the Real Time Mesoscale Analysis (RTMA) and UnRestricted Mesoscale Analysis (URMA) systems at NCEP/EMC             | USA     | 1-25  |
| H. Sako   | Assimilation of JMA high-resolution radiosonde observation data into the mesoscale 4D-Var data assimilation system                         | Japan   | 1-27  |
| H. Seko, T. Yoshihara, A. Senoguchi   | Data assimilation experiment of SSR mode-s downlink data   | Japan   | 1-29  |
| L. Torrìsi, F. Marcucci   | The Italian Air Force Met Service operational NWP system   | Italy   | 1-31  |
| K. Yamashita  | Assimilation of Himawari-8 atmospheric motion vectors into JMA's operational global, mesoscale and local NWP systems                       | Japan   | 1-33  |

## Table of Contents

### 2. Data sets, diagnostic and dynamical investigations, statistical postprocessing, multi-year reanalyses and associated studies

| Author  | Paper Title  | Country        | Pages |
|---|--|----------------|-------|
| M. Akperov, A. Rinke, I. Mokhov, D. Handorf, K. Dethloff and the Arctic Cordex Team | Cyclone activity in the Arctic from an ensemble of regional climate models (Arctic CORDEX)             | Russia/Germany | 2-03  |
| M. G. Akperov, I.I. Mokhov  | Polar lows over Nordic seas from satellite observations and reanalysis data                            | Russia         | 2-05  |
| I. I. Mokhov, M. G. Akperov, M.A Dembitskaya.                                       | Lapse-rate feedback assessment from reanalysis data  | Russia         | 2-07  |
| I. I. Mokhov, A.V. Timazhev   | Climate anomalies and tendencies of change in Lake Baikal basin  | Russia         | 2-09  |
| A.E. Pokhil, I.A. Gorlach   | Blocking synoptic pattern : formation of “small” eddies in the troposphere                             | Russia         | 2-11  |
| A.E. Pokhil   | Numerical simulations of “small” eddies formation in the atmosphere                                    | Russia         | 2-13  |
| A.V. Timazhev   | Frequency of blocking anticyclones in the Northern Hemisphere from RIHMI data: Interannual variability | Russia         | 2-15  |

### 3. Computational studies including new techniques, the effect of varying model resolution, parallel processing

| Author  | Paper Title  | Country | Pages |
|---------|--|---------|-------|
| T. Kato | Influence of horizontal resolution on structure changes of atmospheric stratification in the 2015 Hiroshima heavy rainfall | Japan   | 3-03  |

### 4. Parameterization of important atmospheric and surface processes, effects of different parameterizations

| Author  | Paper Title   | Country | Pages |
|---|---|---------|-------|
| K. Araki  | Influence of Cloud Microphysics Scheme and Ice Nuclei on Forecasting a Heavy Snowfall Event in Japan associated with the “South-Coast Cyclones” | Japan   | 4-03  |
| T. Komori, A. Shimpo  | Improvement of MJO prediction and moistening processes for the DYNAMO period  | Japan   | 4-05  |
| P. Marquet  | The mixed-phase version of moist-air entropy  | France  | 4-07  |
| J. Wang, S. Lu, X. Zhang, S. Kondragunta, A. da Silva, J. McQueen, P. Bhattacharjee, S. Moorthi, A. Harris, P. Koner, Y-T. Hou, V. Tallapragada | The implementation of NEMS GFS Aerosol Component (NGAC) Version 2: Global aerosol forecasting at NCEP using satellite-based smoke emissions     | USA     | 4-09  |
| Y. Yamada   | Development of a two-moment three-ice bulk microphysical model for ice  | Japan   | 4-11  |

## Table of Contents

### 5. Development of and studies with regional and smaller-scale atmospheric models, regional ensemble, monthly and seasonal forecasting

| Author  | Paper Title  | Country   | Pages |
|---|--|-----------|-------|
| L.-P. Caron, C. A. Mehta, J. M. Collins, A-S. Daloz   | Prospects of hurricane variability analysis in dynamically downscaled tropical cyclone simulations           | Spain/USA | 5-03  |
| G.-P. Lou, Q. Liu, E. Rogers, G. DiMego, D. Jovic   | Tropical Storm Relocation in the North America Model (NAM)   | USA       | 5-05  |
| J. McQueen  | NOAA's National Air Quality Forecast Capability for ozone and fine particulate matter                        | USA       | 5-07  |
| J. McQueen  | NWS HYSPLIT atmospheric transport and dispersion modeling  | USA       | 5-08  |
| M. Tsyrlunikov, D.Gayfulin  | A spatio-temporal Stochastic Pattern Generator for use in ensemble prediction and ensemble data assimilation | Russia    | 5-10  |
| V. Tallapragada and the team of the Global Climate and Weather Modeling Branch, Environmental Modeling Center | Upgrade of the NCEP GFS to a 4D Hybrid Ensemble Variational Data Assimilation                                | USA       | 5-12  |
| Y. Wu, M. Ek  | Improve and maintain LSM in NAM  | USA       | 5-14  |

### 6. Developments in global forecast models, case studies, predictability, investigations, global ensemble, monthly and seasonal forecasting

| Author   | Paper Title   | Country | Pages |
|--|---|---------|-------|
| L. Auger, E. Bazile, L. Berre, P. Brousseau, Y. Bouteloup, F. Bouyssel, N. Boullot, P. Cébron, P. Chambon, G. Desroziers, R. El Khatib, G. Faure, C. Fischer, V. Guidard, F. Guillaume, A. Joly, C. Labadie, C. Loo, J.F. Mahfouf, P. Marguinaud, P. Marquet, A. Mary, N. Merlet, L.F. Meunier, Y. Michel, P. Moll, C. Payan, J.M. Piriou, D. Ricard, Y. Seity, F. Taillefer, F. Voitus, E. Wattrelot, K. Yessad | The 2015 upgrades of the Météo-France NWP system  | France  | 6-03  |
| H-M. H. Juang  | Improving spherical harmonic coefficients for the fine resolution NCEP global atmospheric spectral modeling | USA     | 6-05  |
| I.I. Mokhov, A.V. Timazhev   | Summer climate anomalies in the Northern Eurasia during intra-annual transitions from El Niño to La Niña    | Russia  | 6-07  |
| I.I. Mokhov, A.V. Timazhev   | Weather-climate anomalies in Russian regions: El Niño-associated predictability                             | Russia  | 6-09  |

## Table of Contents

### 7. Global and regional climate models, sensitivity and impact experiments, response to external forcing

| Author   | Paper Title  | Country | Pages |
|--|--|---------|-------|
| S. N. Denisov, M.G. Akperov, M. M. Arzhanov, A.V. Eliseev, V.S. Kazantsev, N.V. Pankratova, M.A. Dembitskaya, Yu. A. Shtabkin, E.V. Berezina | Changes in wetland methane emissions in the IAP RAS global model under RCP anthropogenic scenarios                             | Russia  | 7-03  |
| M. Déqué, A. Alias, S. Somot, O. Nuisser   | Climate change and extreme precipitation: the response by a convection-resolving model   | France  | 7-05  |
| I. I. Mokhov, A. V. Timazhev   | Changes of atmospheric blockings in the 21st century from CMIP5 ensemble simulations with RCP scenarios                        | Russia  | 7-07  |
| E. D. Nadyozhina, I. M. Shkolnik, A.V. Sternat, R. S. Bortkovskii  | Spatiotemporal structure of changes in carbon dioxide and oxygen fluxes through the surface of Arctic seas in changing climate | Russia  | 7-09  |
| I. Shkolnik, T. Pavlova, S. Efimov   | Dangerous hydrological events in future climate as projected by an ensemble of high resolution RCM for northern Eurasia        | Russia  | 7-10  |
| N. S. Sidorenkov   | The schedule of the Mul'tanovskii's synoptic periods   | Russia  | 7-12  |

### 8. Development of and advances in ocean modelling and data assimilation, sea-ice modelling, wave modelling

| Author   | Paper Title  | Country | Pages |
|--|--|---------|-------|
| S. Flampouris, J.-H. Alves, M. Pondeva, J. Whiting   | Inclusion of significant wave height analysis to NCEP's UnRestricted Mesoscale Analysis (URMA)                             | USA     | 8-03  |
| Z. D. Garraffo, P. Hogan, A. Mehra, I. Rivin, B. Balasubramanian, T. Spindler, H-C. Kim, S. Paturi | Surface heat flux corrections for Global Ocean Forecasts   | USA     | 8-05  |
| H-C. Kim, A. Mehra, S. Nadiga, Z. Garraffo, S. Son   | Simulated global HYCOM (GLBa0.24) results from various ocean color forcings: Preliminary results from sensitivity analyses | USA     | 8-07  |
| V. Krasnopolsky, S. Nadiga, E. Bayler, A. Mehra, D. Behringer                                      | Consistent ocean color and its assimilation in ocean models  | USA     | 8-09  |
| B. Li, A. Mehra, E. Bayler   | SMOS satellite sea-surface salinity data: impact on upper-ocean modelling  | USA     | 8-11  |
| T. Spindler, A. Mehra, D. Spindler   | Applying multi-model superensemble methods to global ocean operational systems   | USA     | 8-13  |

## 9. Development of and studies with coupled ocean-atmosphere models

| Author                                    | Paper Title  | Country | Pages |
|---|--|---------|-------|
| H-S. Kim, D. Iredell, S. Trahan, A. Mehra | The impact of ocean conditions on the hurricane Blanca (2015) forecasts with a coupled model   | USA     | 9-03  |
| A. Wada, M. Kunii                         | The effect of predicted oceanic conditions on the assimilation of Typhoon Sinlaku (2008)   | Japan   | 9-05  |
| A. Wada                                   | Idealized storm evolution and the difference between the eastern and the western North Pacific calculated by an atmosphere-wave-ocean coupled model                      | Japan   | 9-07  |
| A. Wada                                   | Comparison of numerical simulations of Typhoon Haiyan in 2013 and Typhoon Mike in 1990   | Japan   | 9-09  |
| A. Wada                                   | Typhoon Man-yi in 2013 simulated by an atmosphere-wave-ocean coupled model with 1.2-km horizontal resolution   | Japan   | 9-11  |
| A. Wada                                   | Extremely deepening of central pressures for Typhoon Neoguri in 2014 simulated by an atmosphere-wave-ocean coupled model and its dependency on the horizontal resolution | Japan   | 9-13  |